

Förebyggande mot illamående (Steg 5b, 4d)

Antiemetics

Regimen schedule

Day	1	2	3	4	5	6
Aprepitant Morgondos	X	X	X	X		
Ondansetron Morgondos	X	X	X	X		
Ondansetron Kvällsdos	X	X	X	X		
Betametason Morgondos	X	X	X	X	X	X
Olanzapin As needed						

About your treatment

You must take preventive treatment to reduce the risk of nausea in connection with your treatment. Some medicines also reduce the risk of allergy that certain treatments can cause. If you miss taking your dose, tell your nurse. It is important that you note how you feel so that we can adjust the dose of medicine.

How to take your treatment

Day 1

About 1-2 hours before treatment

Emend (aprepitant) 125mg: 1 capsule

Betapred (betamethasone) 0.5 mg: 12 tablets (tablets dissolve in water)

Ondansetron 8mg: 1 tablet

In the evening

Ondansetron 8mg: 1 tablet

Day 2

About 1-2 hours before treatment

Emend (aprepitant) 80 mg: 1 capsule

Ondansetron 8mg: 1 tablet

Betapred (betamethasone) 0.5 mg: 12 tablets (tablets dissolve in water) **In the evening**

Ondansetron 8mg: 1 tablet

Day 3

About 1-2 hours before treatment

Emend (aprepitant) 80 mg: 1 capsule

Ondansetron 8mg: 1 tablet

Betapred (betamethasone) 0.5 mg: 12 tablets (tablets dissolve in water) **In the evening**

Ondansetron 8mg: 1 tablet

Day 4

About 1-2 hours before treatment

Emend (aprepitant) 80mg: 1 capsule Ondansetron 8mg: 1 tablet Betapred (betamethasone) 0.5 mg: 12 tablets (tablets dissolve in water) **In the evening** Ondansetron 8mg: 1 tablet

Day 5

In the morning

Betapred (betamethasone) 0.5 mg: 8 tablets (tablets dissolve in water)

Day 6

In the morning

Betapred (betamethasone) 0.5 mg: 4 tablets (tablets dissolve in water)

If necessary, treatment days and a few days after

Olanzapine 2.5 mg. If necessary, you can increase to a maximum of four tablets (10 mg) per day. It is best taken at night because the medicine causes fatigue.

Preventive medicine against nausea

You can get medicine from your nurse or by prescription.

Aprepitant

Aprepitant are capsules for nausea. The capsules are taken for three days, starting with a higher dose on the first day. Constipation is common. This can be prevented with laxatives. Inform the healthcare staff before the treatment if you have a sluggish stomach. Common with headache. This can be treated with paracetamol. Dizziness and fatigue may occur. Keep that in mind if you're going to drive. Hiccups also occur.

Ondansetron

Ondansetron are tablets that are swallowed whole with or without food. The dose is taken at least one hour before treatment. Constipation is common from ondansetron. This can be prevented with laxatives. Inform the healthcare staff before treatment if you have a sluggish stomach. Headaches are also common. This can be treated with paracetamol.

Betapred (betametason)

Betapred (betamethasone) is cortisone. The tablets are dissolved in a little water and the solution is swallowed. The dose is taken at least one hour before treatment. Please rinse your mouth with water after taking betamethasone to reduce the risk of oral thrush. Take the tablets with food. Cortisone is also used to reduce the risk of allergic reactions that some cancer drugs can cause. Cortisone can be invigorating and redness.

Planzapin

Olanzapine is a drug used for mental illness, but in low doses has a good effect against nausea, especially against nausea that can occur days after treatment. Olanzapine may make you tired. Fatigue can reduce attention when, for example, driving, which can make it unsuitable to drive a motor vehicle. We recommend taking the dose in the evening, with the first dose the evening of the treatment day. You should take no more than 10 mg of Olanzapine in a day. A higher dose does not give a better effect.

Nausea diary

Register your condition and report back to your nurse.

Name:		Social security number:										
Treatment:												
Course number:							Start date:					
Date of day 1:		Nausea			Vomiting		Appetiteless		Perceived well-being			Notes
		None	Slight	Severe	Yes	No	Yes	No	Good	Poor	Very poor	
Day 1	Evening											
	Morning											
Day 2	Day											
	Evening											
Day 3	Morning											
	Day											
Day 4	Evening											
	Morning											
Day 5	Day											
	Evening											
Day 6	Morning											
	Day											
Day 7	Evening											
	Morning											
Day 8	Day											
	Evening											
Day 9	Morning											
	Day											
Day 10	Evening											
	Morning											