

Melfalan-Prednisolon

for Haematological malignancy

Patient information

Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Melfalan	X	X	X	X																	
Prednisolon	X	X	X	X																	

Next course of treatment, day 36

Day	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Melfalan														
Prednisolon														

About your treatment

Melfalan (Alkeran) is a cytostatic drug. Prednisolone is a cortisone preparation.

Each course is 35 days. Melfalan and prednisone are taken as tablets on days 1-4. After that, there is a break of 31 days, before a new course starts.

During treatment, your blood tests and any side effects will be monitored.

How to take the medicines

Melfalan (Alkeran): Swallow the tablets whole with water half an hour before a meal.

See also information in the package leaflet.

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips: Try to exercise and move around. It's the only thing known to help with fatigue.

Nausea and taste changes

There is a risk that the treatment will make you feel unwell. If necessary, you will be given anti-nausea medication.

Tips: Sometimes it feels better to eat lighter food and small portions. For example, try skim milk, yoghurt, omelettes, sandwiches and hot or cold soups. Drinking a little extra between meals can reduce nausea.

Problems with stomach and intestines

Diarrhoea is common. You may also become constipated.

Tip: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. If you're constipated, try high-fibre foods and exercise. Talk to your doctor if you get diarrhoea or constipation - you may need medication.

Infection susceptibility

You will be susceptible to infection after treatment. This is because the level of white blood cells drops. Before each treatment, blood samples are taken to see if the blood cells have recovered. If the values are too low, the treatment must be postponed.

Tips: Try to avoid close contact with people who have a cold or stomach illness, for example. Wash your hands often with soap and water. Hand sanitizer can be useful.

Problems with skin, hair and nails

Skin often becomes dry and more sensitive to the sun. You may also lose hair or your hair may change texture. Nails can become brittle.

Tip: When washing your skin, be gentle and use mild soap. Lubricate often with moisturiser. Wear covering clothing to protect against the sun, and sunscreen on parts of your body that you can't cover. Wear gloves for work that is dirty or abrasive for your hands. If your nails are bothering you, you can use nail polish and brush your cuticles with oil or apply a moisturiser.

Common advice

Cancer drugs are usually excreted in the urine up to 7 days after treatment. Therefore, you should avoid splashing urine. Sit down when you pee. When finished, close the toilet lid and flush 2 times. It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own. If you are a man, use a condom during sexual intercourse within 72 hours of treatment, as cytostatic drugs are also excreted through the seminal fluid.

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes
