

Regimen schedule

Day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Osimertinib	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

Day	22	23	24	25	26	27	28
Osimertinib	X	X	X	X	X	X	X

About your treatment

Osimertinib (Tagrisso) blocks proteins that regulate the growth and division of cancer cells, slowing it down. The tablets are taken once a day without interruption. During treatment, your blood tests and any side effects will be monitored.

How to take the medicines

Swallow the tablets whole with water. Take the tablets at the same time each day with or without food. If you miss a dose, take it only if there are more than 12 hours before the next dose.

You must not eat or drink St. John's wort preparations during treatment.

See also the information in the package leaflet.

Side effects - symptoms and tips

The side effects we list here are the most common and important. They can vary greatly from person to person, and even from time to time. Side effects may determine your dose or require you to stop taking the medicine.

Tell your doctor or nurse if you have any side effects. Often the symptoms can be prevented or alleviated.

Fatigue

Fatigue is common. You may feel low in energy and never rested. Your memory may deteriorate, and you may find it harder to concentrate and solve problems. It can make you feel down, stressed and anxious. You may also find it difficult to socialise. Fatigue may persist for a long time after treatment.

Tips: Try to exercise and move around. It's the only thing known to help with fatigue.

Acne-like skin problems

You may get acne-like rashes on your face and upper body. You may also experience itching, cracking of the skin and inflammation around the nails. Talk to your nurse if you develop skin problems - you may need medication.

Tip: Apply a moisturising cream to your whole body several times a day. Use bath oil instead of soap. Avoid sun-bathing. Wear a hat and full-coverage clothing. Lubricate parts of the body not covered by clothing with sunscreen that has a high SPF and UVA protection.

Problems with stomach and intestines

Diarrhoea is quite common.

Tip: Drink a lot, a few glasses more than usual per day. If you have diarrhoea, avoid fatty, spicy and high-fibre foods. Talk to your doctor if you get diarrhoea - you may need medication.

Problems with mouth and mucous membranes

The mucous membranes of your nose, mouth, eyes and abdomen may be affected by your treatment. For example, you may have nosebleeds and a runny nose. In the mouth, you may experience dryness, redness, burning and sores. You can also get fungus, which turns the lining of your mouth red or gives it a white coating. Eyes may become dry and watery. The mucous membranes in the lower abdomen can also become dry.

Tip: If you have a dry mouth, use saliva stimulants available in pharmacies. Rinsing your mouth with Vichy water can prevent fungus. Be attentive to oral hygiene, and use a soft toothbrush and mild toothpaste. Be careful with floss and toothpicks. Inform your dentist of your cancer treatment when you visit. Avoid contact lenses if you have eye problems. Women can use oestrogen cream, emollient cream or oil to treat dry mucous membranes in the lower abdomen.

Contact your clinic immediately at:

Acute

Call 112 in case of persistent acute chest pain, pressure on the chest or difficulty breathing/shortness of breath

Call your clinic immediately at:

Fever (temp above 38 degrees) or at a temperature below 36 degrees

Sudden deterioration regardless of symptoms

Severe diarrhea that does not decrease with stopping drugs

New severe fatigue and/or muscle weakness or involuntary muscle cramps

New skin rashes over large parts of the body

Painful skin redness/rash/nail changes

Liquifying rash with yellowish crusts

Painful or inflamed eyes, sensitivity to light, blurred vision or altered vision

Common advice

It is important not to get pregnant or make someone pregnant while you are being treated with cancer drugs, as the drugs can affect the foetus. Therefore, use safe contraceptives. Sometimes double protection is needed, as side effects such as diarrhoea or the direct effect of cancer drugs may mean that the contraceptive pill is not enough protection on its own.

General information about cancer

Useful information about cancer is available from both 1177 Vårdguiden and Cancerfonden:

www.1177.se

www.cancerfonden.se

Notes
